Proximal Humerus Fracture Nonoperative Protocol

	Phase I: Weeks 0-2	Phase II: Weeks 3-6	Phase III: Weeks 6-12	Phase IV: Weeks 12+
Weight-Bearing and Immobilization	Non-weight-bearingSling at all times except as needed to perform PT exercises		Discontinue sling Gradually progress to WBAT	No restrictions
Aerobic	• None	• None	Stationary bike	Begin walk-jog-run progression if desired
Range of Motion (ROM) and Strengthening	 ROM exercises 4-5 times daily ▶ Pendulum exercises ▶ Hand squeezes and pumps ▶ AROM of neck, elbow, wrist, and fingers 	 Continue Phase I exercises Begin supine ER to 15-20° with a slight abduction using a stick or bar Begin gentle forward elevation with pulley assistance At 4 weeks may begin isometric deltoid, IR, and ER exercises at neutral rotation only 	 Continue Phase II exercises Begin light stretching and early active light resistance exercises Initiate active forward elevation while supine (gravity eliminated) and transition to standing with bar assistance and eventually without assistance Bands for progressive and gradual strengthening (flexion, extension, abduction, ER, and IR) 	 Continue Phase III exercises Begin isotonic strength exercises and progress to weights (start at 1 lbs. and progress gradually and as tolerated) Rotator cuff and scapular stretching Progress overhead activities when pain- free and as tolerated
Other	 Heat before and/or ice after PT sessions as needed Modalities as needed to reduce pain/swelling and for muscle re-education Home Exercise Program (HEP) per discretion of Physical Therapist 			
Goals	Pain controlProtect tissue healingMinimize joint stiffness	Pain controlProtect tissue healingBegin ROM exercises	Advance ROM exercises Begin strengthening	 Achieve full ROM Progress activities as tolerated when pain-free